

**ATTACHMENT C.**

---

Wildlife Potential for Occurrence Table

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<b>Amphibians</b>				
<i>Spea hammondi</i> western spadefoot	-	SSC	Prefers open areas with sandy or gravelly soils, in a variety of habitats including mixed woodlands, grasslands, coastal sage scrub, chaparral, sandy washed, lowlands, river floodplains, alluvial fans, playas, alkali flats, foothills, and mountains. Rain pools which do not contain bullfrogs, fish, or crayfish are necessary for breeding.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<b>Birds</b>				
<i>Agelaius tricolor</i> tricolored blackbird	-	ST; SSC	Freshwater marshes with dense cattails, bulrushes, sedges, and tule. Forages in open habitat such as cultivated fields and pastures.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Aimophila ruficeps canescens</i> southern California rufous-crowned sparrow	-	WL	Coastal sage scrub, dominated by CA sagebrush, or in coastal bluff scrub with low scattered scrub and moderate to steep, dry, and rocky slopes. Nests on ground or within 1 meter of ground in shrubs or trees.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Artemisospiza belli belli</i> Bell's sage sparrow	-	WL	Chaparral dominated with California sagebrush or chamise. Nests on ground or within 1 meter above ground in a shrub.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Athene cunicularia</i> Burrowing owl	-	SSC	Open grasslands including prairies, plains, and savannah, or vacant lots and airports. Nests in abandoned dirt burrows.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Buteo swainsoni</i> Swainson's hawk	-	ST	Open pine-oak woodland, savannah, and agricultural fields with scattered trees. Nests in solitary bush or tree, or in small groves.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<i>Coccyzus americanus occidentalis</i> western yellow-billed cuckoo	FT	FE	Open woodland habitat, near water, especially with dense willow and cottonwood understory.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Eremophila alpestris actia</i> California horned lark	-	WL	Bare open areas dominated by low vegetation or widely scattered shrubs, includes prairies, deserts, and plowed fields. Nests in a hollow on the ground.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Falco columbarius</i> merlin	-	WL	Open woodland and fragmented forests. Also found in grasslands, marshes, deserts, lakes, fields, along the coast.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Laterallus jamaicensis coturniculus</i> California black rail	-	ST; FP	Coastal and estuarine saltmarshes especially dominated by pickleweed and matted salt grass. Freshwater marshes with shallow and stable water levels and flat shorelines.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Polioptila californica californica</i> coastal California gnatcatcher	FT	SSC	Dry coastal slopes, washes, and mesas with areas of low vegetation and coastal sage scrub.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Vireo bellii pusillus</i> least Bell's vireo	FE	SE	Riparian woodlands and willow-cottonwood forests particularly with streamside thickets and dense brush.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<b>Fish</b>				
<i>Catostomus santaanae</i> Santa Ana sucker	FT	-	Pools and runs of creeks and small to medium rivers with cool, shallow, clear, and unpolluted water.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<i>Gila orcuttii</i> arroyo chub	-	SSC	Creeks, streams, and rivers with areas of slow-moving water with sand or mud bottoms. Ranges from San Diego to San Luis Obispo county.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Oncorhynchus mykiss irideus pop. 10</i> steelhead - southern California DPS	FE	-	Freshwater lagoons, river mouths/tidal rivers, bays/sounds, rivers, and creeks for breeding. Spawning streams are relatively warm. Migrate between freshwater breeding and marine non-breeding habitats. Marine habitat is pelagic and near shore.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<b>Insects</b>				
<i>Bombus crotchii</i> Crotch bumble bee	-	SC	Grasslands and shrublands in a hotter and drier environment than other bumblebee species. Nests underground, often in abandoned rodent dens. Most observations occur in southern California in coastal areas.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Eugnosta busckana</i> Busck's gallmoth	-	-	Coastal sand dunes.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Euphydryas editha quino</i> quino checkerspot butterfly	FE	-	Chaparral and coastal sage scrublands in Riverside and San Diego counties.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Rhaphiomidas terminatus abdominalis</i> Delhi Sands flower-loving fly	FE	-	Dune habitat, with fine sandy Delhi soils.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<b>Mammals</b>				

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<i>Chaetodipus fallax fallax</i> northwestern San Diego pocket mouse	-	SSC	Coastal scrub, chaparral, sagebrush, and grasslands in western San Diego county.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Dipodomys merriami parvus</i> San Bernardino kangaroo rat	FE	SC; SSC	Alluvial sage scrub, flood plains, washes, and upland areas adjacent to desert habitat.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Dipodomys stephensi</i> Stephens' kangaroo rat	FE	FT	Annual grasslands, coastal sage scrub with sparsely spaced vegetation, loose friable soils, and flat or slightly rolling terrain.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Eumops perotis californicus</i> western mastiff bat	-	SSC	Roosts high above ground in rock and cliff crevices, shallow caves, and rarely in buildings. Occurs in arid and semiarid regions including rocky canyon habitats.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Lasiurus xanthinus</i> western yellow bat	-	SSC	Roosts in trees, especially in fan palms with dead fronds. Found in riparian woodlands in arid regions, oak or pinyon-juniper woodlands, and human developed areas.	<b>Moderate:</b> Habitat for the species occurs on site and a historic known occurrence has been recorded within five miles.
<i>Lepus californicus bennettii</i> San Diego black-tailed jackrabbit	-	SSC	Variety of open or semi-open country including grasslands, croplands, and sparse coastal scrub.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Nyctinomops femorosaccus</i> pocketed free-tailed bat	-	SSC	Roosts high above ground in rock and cliff crevices, shallow caves, and rarely in buildings. Occurs in arid and semiarid regions including rocky canyon habitats.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<i>Onychomys torridus ramona</i> southern grasshopper mouse	-	SSC	Low, semi-open, and open scrub habitats with flat, sandy valley floors. Habitats include coastal and mixed chaparral, coastal sage scrub, riparian scrub, low sagebrush, and grasslands with interspaced shrubs.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Perognathus longimembris brevinasus</i> Los Angeles pocket mouse	-	SSC	Habitats with sandy and fine soils, including grasslands, coastal sage scrub, and alluvial sage scrub.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<i>Taxidea taxus</i> American badger	-	SSC	Open habitats with friable soil such as grasslands, brushlands with sparse ground cover, open chaparral, and sometimes riparian zones.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.
<b>Reptiles</b>				
<i>Anniella stebbinsi</i> southern California legless lizard	-	SSC	Occurs in moist warm loose soil with plant cover. Can occur in sparsely vegetated areas of coastal sand dunes, chaparral, pine-oak woodlands, desert scrub, sandy washes, alluvial fans, and stream terraces with sycamores, cottonwoods, or oaks.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Arizona elegans occidentalis</i> California glossy snake	-	SSC	Arid scrub, rocky washes, grasslands, and chaparral. Prefers microhabitats of open areas and areas with soil loose enough for easy burrowing.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Aspidoscelis hyperythra</i> orange-throated whiptail	-	WL	Semi-arid open areas with coarse soils including coastal sage scrub, chaparral, and dry riparian areas and washes.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.

Wildlife Species Occurrence Table				
Species	Status		Habitat	Potential for Occurrence
	Fed	CA		
<i>Aspidoscelis tigris stejnegeri</i> coastal whiptail	-	SSC	Arid habitats including chaparral, woodlands, and dry riparian areas. Primarily hot and dry open areas with sparse foliage.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Coleonyx variegatus abbotti</i> San Diego banded gecko	-	SSC	Rocky areas in coastal sage scrub and chaparral.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Crotalus ruber</i> red-diamond rattlesnake	-	SSC	Found in coastal chaparral, arid scrub, rocky grassland, oak and pine woodlands, desert mountain slopes and rocky desert flats.	<b>Presumed absent:</b> No suitable habitat is present on site; known occurrence has been recorded within five miles.
<i>Phrynosoma blainvillii</i> coast horned lizard	-	SSC	Open areas of valleys, foothills, and semiarid mountains with sandy soil and low vegetation including chaparral, woodlands, and grasslands.	<b>Presumed absent:</b> No suitable habitat is present on site. Historical known occurrence has been recorded within five miles.

**U.S. Fish and Wildlife Service (USFWS) – Federal**

FE Federal Endangered  
FT Federal Threatened

**California Department of Fish and Wildlife (CDFW) – California**

SE State Endangered                      SSC Species of Special Concern  
ST State Threatened                      FP California Fully Protected  
SC State Candidate for Listing        WL Watch List